

Azimuthal and pseudo-rapidity correlations with strange particles at intermediate- p_T at RHIC

J Bielcikova for the STAR Collaboration

Physics Department, Yale University, New Haven, CT-06520, USA

E-mail: jana.bielcikova@yale.edu

Abstract. We present results on two-particle azimuthal correlations with strange trigger particles (K_S^0 , Λ , Ξ , Ω) associated with unidentified charged particles in d+Au and Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV. We investigate, in detail, the near-side associated yield as a function of centrality, p_T and strangeness content in the trigger particle to look for possible flavor and baryon/meson differences. We compare our results to a fragmentation and recombination model, where the study of Ω -triggered correlations is used as a critical test of the validity of the recombination picture.

PACS numbers: 25.75.-q, 25.75.Gz

1. Introduction

The suppression of inclusive p_T spectra of identified particles in central Au+Au collisions with respect to p+p and peripheral Au+Au collisions [1, 2] and the enhanced baryon/meson ratios [3, 4] show that baryons and mesons behave differently than in p+p collisions for $p_T = 2$ -6 GeV/c. This indicates that fragmentation is not dominant and parton recombination and coalescence models [5, 6, 7, 8] have been suggested as alternative mechanisms of particle production. Moreover, studies of di-hadron correlations in Au+Au revealed the presence of an additional long-range pseudo-rapidity correlation (*ridge*) on the near-side [9], which is absent in p+p and d+Au collisions.

In this paper, we discuss the properties of two-particle correlations using strange trigger particles (K_S^0 , Λ , Ξ , Ω) associated with unidentified charged particles in d+Au and Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV with STAR. We investigate the near-side associated yield as a function of centrality, system size, p_T and strangeness content in the trigger particle, to look for possible flavor and baryon/meson differences. We compare our results to a fragmentation and recombination model.

2. Data analysis

The correlation functions, normalized to the number of trigger particles, are corrected for the reconstruction efficiency of associated particles, elliptic flow (v_2) and, unless mentioned otherwise, triangular acceptance in $\Delta\eta$. The near-side yield of associated




Figure 1. Centrality dependence of the ridge yield (a) and jet yield (b) of associated charged particles for various trigger species in d+Au and Au+Au collisions. The error bands indicate systematic errors on the ridge yield due to the v_2 subtraction.

particles is calculated as the area under the Gaussian peak obtained from a fit. We study separately the jet and ridge contributions to the near-side yield by analyzing the correlations in two $\Delta\eta$ windows: $|\Delta\eta| < 0.7$ containing both jet and ridge correlations, and $|\Delta\eta| > 0.7$ containing only the ridge contributions, assuming the jet contribution at large $\Delta\eta$ is negligible. The jet yield is free of systematic uncertainties due to the v_2 subtraction if a uniformity of v_2 with η is assumed. For the ridge yield, these systematic errors are estimated by subtracting the v_2 measured by the event plane method (the lower bound) and by the 4-particle cumulant method (the upper bound).

3. Correlations with K_S^0 , Λ and charged trigger particles

Comparing the near-side yields in d+Au and Au+Au collisions, we observe a strong increase by a factor of 3-4 going from d+Au to central Au+Au collisions. Studying separately the jet and ridge contributions to the near-side yield (Fig. 1), we find that the ridge yield rises with centrality and is responsible for the observed strong increase of the near-side yield with centrality. The jet yield is independent of centrality and consistent with that in d+Au collisions. No significant baryon/meson or particle/anti-particle differences are observed.

Next, we study the dependence of the near-side yield on the transverse momentum of the trigger particle, p_T^{trig} , shown in Fig. 2. While the ridge yield increases with p_T^{trig} and flattens off for $p_T^{trig} > 3.0$ GeV/c, the jet yield keeps increasing with p_T^{trig} as expected. The jet yield for Λ triggers is systematically below that of charged hadron and K_S^0 triggers. Remaining effects of artificial track merging, which are found to affect more V0s than charged tracks, are currently under investigation.

We have also measured the invariant p_T spectra of associated charged particles for




Figure 2. Dependence of the ridge yield (a) and jet yield (b) on p_T^{trig} for various trigger species in central (0-10%) Au+Au collisions. The bands indicate systematic errors on the ridge yield due to the v_2 subtraction.

$p_T^{trig} = 3-6$ GeV/c (not shown) and extracted the inverse slope, T . The ridge spectra have, for all studied trigger species, $T \sim 400$ MeV, close to that of 'the bulk', while the jet spectra have $T \sim 450$ MeV. The 0-10%/40-80% centrality ratio of the near-side yields is about 3 at $p_T^{assoc} = 1$ GeV/c and decreases with p_T^{assoc} . As shown in Fig. 1, this large ratio is due to the correlations at large $|\Delta\eta|$ because the jet yield is independent of centrality. Our results qualitatively agree with the parton recombination model [10] which points toward a significant role of thermal-shower recombination in Au+Au collisions. To draw quantitative conclusions, the calculation must be done for the same centrality and p_T^{trig} selection and also reproduce properties of the measured $\Delta\eta$ correlations.

4. Correlations with multiply-strange trigger baryons

Our study of correlations with multiply-strange baryons, especially Ω , has been stimulated by the predictions from the recombination model of [11]. Contrary to the production of K and Λ , for particles created exclusively from strange quarks, such as ϕ and Ω , the contribution from shower s quarks should be negligible for p_T up to 8 GeV/c. Consequently, there should be no associated particles for ϕ - and Ω -triggered correlations with $p_T > 3$ GeV/c because they cannot be distinguished from background.

Figure 3(a) shows the azimuthal correlations for strange trigger baryons with increasing strangeness content: Λ , Ξ and Ω . Due to limited statistics, the trigger particles have been selected with $p_T^{trig} = 2.5-4.5$ GeV/c. Clearly, there is a remaining near-side peak above the elliptic flow contribution for all discussed trigger species. The near-side peak persists even if the larger v_2 , determined from the event plane method, is used. Moreover, the strength of the near-side peak is, within errors, independent of the strangeness content in the trigger particle. This is further confirmed by the study of




Figure 3. (a) Azimuthal correlations with Λ , Ξ and Ω for $p_T^{\text{trig}}=2.5\text{--}4.5$ GeV/ c in 0-10% central Au+Au collisions. The lines indicate the v_2 contributions. (b) Dependence of the near-side yield on p_T^{trig} for various strange trigger particles in central (0-10%) Au+Au collisions. The bands represent systematic errors due to the v_2 subtraction. No $\Delta\eta$ acceptance correction has been applied.

the p_T^{trig} dependence of the near-side yield shown in Fig. 3(b). Within errors, the yield is consistent for all studied baryon and meson species.

5. Conclusions

We have reported results on two-particle correlations with strange trigger particles at intermediate- p_T at RHIC. The correlations reveal a strong contribution from the long-range $\Delta\eta$ correlations in the near-side. We do not observe any significant baryon/meson differences. The behavior of the central-to-peripheral ratio of the near-side yields agrees qualitatively with a recombination model. However, for the Ω -triggered correlations there exist associated particles in the near-side, which disagrees with the model. Studies of jet and ridge yields for Ω -triggered correlations as well as correlations with identified associated particles are expected to constrain the origin of these correlations.

References

- [1] Adler S S *et al* (PHENIX) 2003 *Phys. Rev. Lett.* 91 172301
- [2] Adams J *et al* (STAR) 2004 *Phys. Rev. Lett.* 92 052302
- [3] Adams J *et al* (STAR) nucl-ex/0601042
- [4] Abelev B *et al.* (STAR) 2006 *Phys. Rev. Lett.* 97 152301
- [5] Fries R J, Muller B, Nonaka C and Bass S A 2003 *Phys. Rev.* C68 044902
- [6] Greco V, Ko C M and Levai P 2003 *Phys. Rev.* C68 034904
- [7] Greco V, Ko C M and Levai P 2003 *Phys. Rev. Lett.* 90 202302
- [8] Hwa R C and Yang C B 2003 *Phys. Rev.* C67 034902
- [9] Putschke J (STAR) these proceedings
- [10] Hwa R C and Tan Z 2005 *Phys. Rev.* C72 057902
- [11] Hwa R and Yang C B 2006 nucl-th/0602024